BMC Medicine

Developing an integrated microsimulation model for the impact of fiscal policies on child health in Europe: the example of childhood obesity in Italy --Manuscript Draft--

| Manuscript Number: | BMED-D-21-01486R2 | | | | | |
|-----------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|--|
| Full Title: | Developing an integrated microsimulation model for the impact of fiscal policies on child health in Europe: the example of childhood obesity in Italy | | | | | |
| Article Type: | Research article | | | | | |
| Section/Category: | Global, Public and Environmental Health | | | | | |
| Funding Information: | horizon 2020 framework programme (733206) | Not applicable | | | | |
| | horizon 2020 framework programme (774548) | Not applicable | | | | |
| Abstract: | tax-benefit microsimulation program EURO data. Secondly, module 2(M2) exploits data study, and runs a series of concatenated reprospective effects of income on child Body module 3(M3) uses dynamic microsimulatic structure and incomes obtained by M1, with estimated effect sizes provided by M2, projesimulated policy scenarios. Results. Both universal benefits, such as Linterventions, such as Child Benefit (CB) for effect on childhood overweight, with a Previn comparison with the baseline fiscal syste (95%CI:0.83-0.94), respectively. The impact child obesity, reaching PR of 0.67 (95%CI:0.95%CI:0.44-0.84) for CB at the same age. effects, the estimated costs for 1% prevaler respect to the baseline scenario is much low with universal ones. Conclusions. Our results show that fiscal prohildhood health conditions. Focalised interventions. | ol for analysing the prospective effects of ean countries. The aim of this first ernative fiscal policies on childhood be integrated modules. Firstly, module on disposable household income using the MOD fed with the Italian EU-SILC 2010 of provided by the Italian NINFEA birth cohort gressions in order to estimate the Mass Index(BMI) at different ages. Finally, on techniques that combine the population of regression model specifications and ecting BMI distributions according to the Iniversal Basic Income (BI), and targeted or poorer households, have a significant alence Ratio (PR) in 10-years-old children – m - of 0.88 (95%CI:0.82-0.93) and 0.89 of the fiscal reforms was even larger for 0.50-0.83) for the simulated BI and 0.64 While both types of policies show similar not reduction in overweight and obesity with over with more focalised benefit policy than olicies can have a strong impact on ventions that increase family income, as, can help to prevent child overweight and forecast the effects of fiscal policies on | | | | |
| Corresponding Author: | Davide Rasella, PhD ISGLOBAL: Instituto de Salud Global de Ba Barcelona, Catalunya SPAIN | ırcelona | | | | |
| Corresponding Author Secondary Information: | | | | | | |
| Corresponding Author's Institution: | ISGLOBAL: Instituto de Salud Global de Ba | ırcelona | | | | |
| Corresponding Author's Secondary Institution: | | | | | | |
| First Author: | Davide Rasella, PhD | | | | | |
| First Author Secondary Information: | | | | | | |

| Order of Authors: | Davide Rasella, PhD |
|-----------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| | Lorenzo Richiardi, PhD |
| | Nicolai Brachowicz, MSc. |
| | H. Xavier Jara, PhD |
| | Mark Hanson, PhD |
| | Delia Boccia, PhD |
| | Matteo G. Richiardi, PhD |
| | Costanza Pizzi, PhD |
| Order of Authors Secondary Information: | |
| Response to Reviewers: | Editor's Comments: |
| | We require confirmation of authorship from all co-authors, and have not yet received a response from the listed authors. I have resent them the verification email containing the confirmation link. May I kindly ask that you follow up on this or provide us with alternative email addresses, if necessary? Co-authors should contact us directly to confirm co-authorship if they have any problems. Still to confirm co-authorship Delia Boccia Matteo G. Richiardi A: Both co-authors should have confirmed the coauthorship. |
| | Please include a figure legend. Figure legends should be no more than 300 words (advised but not mandatory). A: Now, all figures have the corresponding legend. Section Methods, Figure 1, lines 111-113, page 6, and lines 344-347 page 19. |
| | 3. Each file (whether main or Additional) should be mentioned in the main manuscript text in order. A: Now, all files are mentioned in the main manuscript text in order. Section Methods, lines 104, page 5; line 125, page 6; and line 138, page 7. |
| | 4. Please rename supplementary files as 'Additional file X', and cite explicitly by additional file name in the manuscript e.g. 'Additional file 1: Fig. S1'. Please ensure that if you have more than one additional file that they are cited in ascending order within the main body of text. A: Supplementary files were renamed in accordance with BMC Medicine guidelines. Supplementary information Section, lines 530, 532, 534, page 29. |
| | 5. Please add a description of your additional file to the end of the manuscript. Legends to Combined Additional files should include a short description of the constituent parts, e.g. "Additional File 1: Figures S1-S3. FigS1- [short title]. FigS2 –[short title]. FigS3 – [short title]".Please note that legends are advised to be no more than 300 words, though this is not mandatory. A: Description of additional files added in Supplementary information Section, lines 531, 533, 535, page 29. |
| | 6. Please could you update the reference formatting to remove the extra details of |

Month of publication, date cited, and "Internet". References should follow the format: Smith JJ. The world of science. Am J Sci. 1999;36:234-5. Journal titles should be abbreviated as in the National Library of Medicine https://www.ncbi.nlm.nih.gov/nlmcatalog/journals

A: Thank you for noticing that, we have changed the references style in accordance with BMC Medicine guidelines.

7. Please move all references to the main reference list (i.e. no additional references in Additional Files). These should be included in the main text when the Additional File is first introduced, for example "...see Additional File 1: Table S1) [XX-XX]"; please ensure the reference list is numbered in the order as they appear in the text.

A: References that were previously included in Supplementary material but not were mentioned in the main manuscript are now cited in Section Methods, lines 104-105, page 5

8. Please make the following change to your section headings: change 'Introduction' to 'Background'.

A: We did it. Background Section, line 64, page 4.

Health (MICH) along the text and in the additional files.

9. Manuscripts should have the following subsections: Background/Methods/Results/Discussion/Conclusions. Please add a new subsection entitled Conclusions.

A: We added Conclusions subtitle before the already existing conclusion paragraph. Conclusions Section, line 431, page 24.

10. Please provide a list of all the abbreviations used in the manuscript. Please place this list just before the Declarations section. If abbreviations are used in the text, they should be defined in the text at first use and included in this list.

A: List of Abbreviations was added just before the Declarations section, lines 442-484, pages 24-26. For the sake of simplicity and clarity, the Microsimulation for Income and Child Health Outcomes (MICHO) was renamed Microsimulation for Income and Child

- 11. You should include a "Declarations" section, consisting of the following subsections: Acknowledgments; Consent for Publication; Funding; Availability of Data and Materials; Authors' Contributions; Competing interests; Ethics Approval and Consent to Participate. All of these subsections MUST be present.
- The "Authors' Contributions" section of the Declarations should include the text "All authors read and approved the final manuscript"
- The "Competing Interests" section of the Declarations should include any relevant competing interests, or state "The authors declare that they have no competing interests". See here for our editorial policies and a full explanation of competing interests. https://www.biomedcentral.com/getpublished/editorial-policies#CompetingInterests

A: As requested we added the Declarations Section, with all subsections present, from line 489 of page 26.

Reviewer reports:

Reviewer #1: This is a revised manuscript describing a microsimulation model testing

the impact of a range of fiscal policies on child weight in Italy. The authors have done a good job of addressing the issues raised by myself and the other reviewer. I just have two remaining comments.

The first relates to a comment in the response to reviewers: "To the best of our knowledge there are no randomized trials of fiscal interventions or increased total family disposable income with child BMI as the outcome. There are several quasi experimental studies on cash transfer, but those studies were conducted in a different population (United States) and did not include child BMI as one of the health outcomes." There are actually a handful of quasi experimental studies on cash transfers with BMI as the outcome, across several countries, and these have found mixed results - see review article: Cooper K, Stewart K. (2017) Does money affect children's outcomes? An update. Case paper 203; ISSN 1460-5023. Centre for Analysis of Social Exclusion, London.

ttp://eprints.lse.ac.uk/103494/1/casepaper203.pdf. This should be acknowledged in light of the strong effects implied by the microsimulation.

A: We thank the reviewer for bringing to our attention the review from Cooper et al. In their work they only mention one study about EITC and obesity from Chia et al. 2013. which admittedly we did not know. As suggested by the reviewer, there are indeed a handful of studies on the effect of income support interventions and obesity among children; however, conclusions from these studies need to be interpreted cautiously, because: a. Only few come from high income countries (the vast majority are from middle income countries); b. results can be conflicting as in some cases these interventions have been associated with an increase of obesity among children. This seems to be dependent on the baseline income of the intervention recipients, the age of children at the time of transfer and also the context where the intervention is delivered). Interestingly, in LMICS results become more consistent in the direction of a positive impact when interventions are delivered in combination with health education and in context where food quality and availability is improved. Overall, these findings do not undermine the role of income-support interventions in addressing child obesity, but suggest that their maximum potential is achieved when delivered in combination with health education and within a food secure environment (in terms of food quality). We have added at page 22 line 391: "To the best of our knowledge there are no randomized trials of fiscal interventions or increased total family disposable income with child BMI as the outcome, but a few quasi-experimental studies have evaluated the impact of cash transfers on child obesity. [37] The overall impact of these interventions seems to be dependent on the baseline income of the intervention recipients, the age of children at the time of transfer and also the context where the intervention is delivered [38-40]."

Second, there was another comment in the response to reviewers, "response rates and attrition are at the standard level (after 10 years a lost of around 30% of the participants). Due to that reason, and that the attrition was not differential due to the overall characteristics of the remaining individuals, we didn't use any approach to correct for that." The table in the Supplementary Methods described the level of attrition - thank you - but I couldn't see evidence that the attrition was not differential, though perhaps I missed it. Can this be made clear and a brief comment included in the paper indicating that attrition was not differential?

A: We thank the reviewer for this comment. The sentence about non-differentiality was based on the comparison of the distribution of the weight measures at 18 months between those lost to the 4-year follow-up and those still in the study at that time: mean weight at 18 months in the two groups equal to 11130 and 11191 grams respectively (p-value from T-Student test=0.23). However, we agree with the reviewer that the statement about non-differentiability is too strong as several other factors might affect attrition, so we did not included such statement in the main text or in the additional files, and we have left in the limitations section (p.24 line 433):" We also acknowledge that the NINFEA cohort has attrition, even if within the standard magnitude for such kind of child cohorts."

Editorial Policies

Please read the following information and revise your manuscript as necessary. If your manuscript does not adhere to our editorial requirements, this may cause a delay while this is addressed. Failure to adhere to our policies may result in rejection of your manuscript.

In accordance with BioMed Central editorial policies and formatting guidelines, all manuscript submissions to BMC Medicine must contain a Declarations section which includes the mandatory sub-sections listed below.

Please refer to the journal's Submission Guidelines web page for information regarding the criteria for each sub-section (https://bmcmedicine.biomedcentral.com/).

Where a mandatory Declarations section is not relevant to your study design or article type, please write 'Not applicable' in these sections.

For the 'Availability of data and materials' section, please provide information about where the data supporting your findings can be found.

We encourage authors to deposit their datasets in publicly available repositories (where available and appropriate), or to be presented within the manuscript and/or additional supporting files.

Please note that identifying/confidential patient data should not be shared.

Authors who do not wish to share their data must confirm this under this sub-heading and also provide their reasons.

For further guidance on how to format this section, please refer to BioMed Central's editorial policies page (see links below).

Declarations

|-

Ethics approval and consent to participate

|-

Consent to publish

-

Availability of data and materials

-

Competing interests

-

Funding

|-

Authors' Contributions

ŀ

Acknowledgements

Further information about our editorial policies can be found at the following links: Ethical approval and consent:

http://www.biomedcentral.com/about/editorialpolicies#Ethics

Availability of data and materials section:

http://www.biomedcentral.com/submissions/editorial-

policies#availability+of+data+and+materials

Our flexible approach during the COVID-19 pandemic

If you need more time at any stage of the peer-review process, please do let us know. While our systems will continue to remind you of the original timelines, we aim to be as flexible as possible during the current pandemic.

This letter contains confidential information, is for your own use, and should not be forwarded to third parties.

Recipients of this email are registered users within the Editorial Manager database for this journal. We will keep your information on file to use in the process of submitting, evaluating and publishing a manuscript. For more information on how we use your personal details please see our privacy policy at

https://www.springernature.com/production-privacy-policy. If you no longer wish to receive messages from this journal or you have questions regarding database management, please contact the Publication Office at the link below.

Email: davide.rasella@isglobal.org

Phone: +34679590927

Developing an integrated microsimulation model for the impact of fiscal policies on child health in Europe: the example of childhood obesity in Italy Davide Rasella, PhD ^{1,2}; Lorenzo Richiardi, PhD ²; Nicolai Brachowicz, MSc ¹, H. Xavier Jara, PhD ³; Mark Hanson, PhD ⁴; Delia Boccia, PhD ^{2,5}; Matteo G. Richiardi, PhD ³; Costanza Pizzi, PhD² ¹ ISGlobal, Hospital Clínic - Universitat de Barcelona, Barcelona, Spain ² Department of Medical Sciences, University of Turin, Turin, Italy ³ Centre for Microsimulation and Policy Analysis, Institute for Social and Economic Research, University of Essex, Colchester, UK ⁴ Institute of Developmental Sciences and NIHR Biomedical Research Centre, University of Southampton and University Hospital Southampton, UK. ⁵ Faculty of Population and Health Policy, London School of Hygiene and Tropical Medicine, London, United Kingdom **Corresponding Author:** Davide Rasella, ISGlobal, Hospital Clínic - Universitat de Barcelona, Barcelona, Spain Hospital Clínic - Universitat de Barcelona, Carrer Rosselló 132, Barcelona 08036, Spain

ABSTRACT

Background. We developed an integrated model called Microsimulation for Income and Child Health (MICH) that provides a tool for analysing the prospective effects of fiscal policies on childhood health in European countries. The aim of this first MICH study is to evaluate the impact of alternative fiscal policies on childhood overweight and obesity in Italy.

Methods. MICH model is composed by three integrated modules. Firstly, module 1 (M1) simulates the effects of fiscal policies on disposable household income using the tax-benefit microsimulation program EUROMOD fed with the Italian EU-SILC 2010 data. Secondly, module 2 (M2) exploits data provided by the Italian birth cohort called Nascita e Infanzia: gli Effetti dell'Ambiente (NINFEA), translated as Birth and Childhood: The Effects of the Environment) study, and runs a series of concatenated regressions in order to estimate the prospective effects of income on child Body Mass Index (BMI) at different ages. Finally, module 3 (M3) uses dynamic microsimulation techniques that combine the population structure and incomes obtained by M1, with regression model specifications and estimated effect sizes provided by M2, projecting BMI distributions according to the simulated policy scenarios.

Results. Both universal benefits, such as Universal Basic Income (BI), and targeted interventions, such as Child Benefit (CB) for poorer households, have a significant effect on childhood overweight, with a Prevalence Ratio (PR) in 10-years-old children - in comparison with the baseline fiscal system - of 0.88 (95%CI:0.82-0.93) and 0.89 (95%CI:0.83-0.94), respectively. The impact of the fiscal reforms was even larger for child obesity, reaching a PR of 0.67 (95%CI:0.50-0.83) for the simulated BI and 0.64 (95%CI:0.44-0.84) for CB at the same age. While both types of policies show similar effects, the estimated costs for 1% prevalence

 reduction in overweight and obesity with respect to the baseline scenario is much lower with more focalised benefit policy than with universal ones. Conclusions. Our results show that fiscal policies can have a strong impact on childhood health conditions. Focalised interventions that increase family income, especially in the most vulnerable populations, can help to prevent child overweight and obesity. Robust microsimulation models to forecast the effects of fiscal policies on health should be considered as one of the instruments to reach the Health in All Policies goals (HiAP). Keywords: Fiscal Policies; Poverty alleviation; Microsimulation; Child Health; Child Overweight; Child Obesity.

BACKGROUND

Fiscal policies, including fiscal benefits and tax reductions, are interventions that can quickly and effectively change income of poor households. Literature points to family income, considered as a general indicator of socio-economic position (SEP) [1], as one of the strongest socio-economic determinant of health [2, 3]. Increasing household income, especially among the most vulnerable families, could prevent several health outcomes such as overweight and obesity, that represent a serious public health concern in high-income countries and increasingly in low and middle income countries [4].

On the one hand, some studies have shown strong negative associations between household income and child obesity [5–7] suggesting that implementing income subsidies, especially

income and child obesity [5–7], suggesting that implementing income subsidies, especially among those families that belong to the first deciles of the income distribution, could relax their economic constraints and free resources that may be spent on improving their dietary intake and adopting healthier physical activities. On the other hand, although several mathematical models for obesity reduction have been developed, they all focus on postnatal interventions on diet, physical activity and other lifestyle-related practices, ignoring income and socio-economic position factors [8–10].

The aim of our study is twofold. Firstly, we developed a flexible three-part integrated microsimulation model as a useful policy design tool for investigating the prospective effects of poverty alleviation fiscal policies on child health outcomes, in line with the Health in All Policies (HiAP) framework [11]. Secondly, we provided a case study for Italy that evaluates the potential effects of eight different simulated fiscal policies on overweight and child obesity. The simulated policies range from basic income policies that ensure a universal yearly basic income, to more targeted policies such as monthly child benefits to low income families

with children under 5 years of age. These fiscal policies were chosen to allow for identifying potential dose responses. Our aim is to contribute to the decision-making process by offering an integrated approach that permits to evaluate prospective costs and effects of several fiscal policy interventions on child health.

METHODS


Study design

The modelling strategy involves two phases and three integrated modules. The first phase comprises two modules. Firstly, we simulate the prospective effects of a variation in benefit policies on equivalised income using EUROMOD (Module 1 - M1). Secondly, we run a series of concatenated regressions to estimate parameters of interest of several relationships using NINFEA variables (Module 2 - M2). In the second phase we apply the estimated parameters obtained from M2 to the simulated population obtained from M1 in order to get a simulated BMI distribution (Module 3 – M3).

The overall structure of our Microsimulation for Income and Child Health (MICH) model, and flow of inputs and outputs for each stage are shown in Figure 1. A more detailed description of each phase and module is provided in the following paragraphs. Further information on the modelling process and its parameters is provided in Additional file 1 [12-15], in accordance with international reporting guidelines recommended by the International Society for Pharmacoeconomics and Outcomes Research and the Society for Medical Decision Making (ISPOR-SMDM) [16].

Figure 1 Structure of the MICH Model: flow of input, output data, and parameters

between M1, M2 and M3 modules



Note: For M1: data provided by the Italian EU-SILC 2010 survey. Fiscal Policies simulated using EUROMOD software. For M2: data provided by NINFEA cohort. Concatenated regressions run in Stata. For M3: dynamic microsimulation using R.

Data sources

EUROMOD is a tax-benefit microsimulation model for the European Union (EU) and the United Kingdom (UK). It is a software that allows to compute the effects of taxes and benefits on household incomes and work incentives for the population of each country and for the EU as a whole in a standardized and comparable manner [17].

EUROMOD covers the 28 member states and is updated to recent policy systems using data from the European Union Statistics on Income and Living Conditions (EU-SILC) as the input database [18], supported by Directorate-General for Employment, Social Affairs and Inclusion (DG EMPL) of the European Commission [19][20]. A more detailed description of EUROMOD is provided in the Additional file 2 [21].

We use the Italian EU-SILC 2010 data as the input population for EUROMOD, in order to be consistent with the income values of Italian NINFEA 2011 cohort, given that Italian EU-SILC 2011 data for EUROMOD was not available. Baseline scenario is based on the tax-benefit system corresponding to June 2018, given that this was the most recent system when this study was carried out.

The Nascita e Infanzia: gli Effetti dell'Ambiente (NINFEA, translated as Birth and Childhood: The Effects of the Environment) project [22, 23], is an Italian birth cohort study that aims at investigating the effect of several exposures acting during pre-natal and early post-natal life on later health. Individuals of the cohorts are children whose mothers voluntarily accepted to participate and had enough knowledge of the Italian language to complete the online questionnaires. The first baseline questionnaire on general health and exposures was fulfilled before and during pregnancy. A more detailed description of NINFEA is provided in the Additional file 3 [24]. We used NINFEA database version 02.2019. This database consists of 6625 mothers and 7423 pregnancies. Data on demographic and socioeconomic factors of each household were collected using the baseline questionnaire completed during pregnancy. Using these aforementioned variables, namely parental age, cohabitation status, education, country of birth, occupation, house size and type, and family size, and external data from the Italian EU-SILC 2011 survey, the Equivalised Household Income Indicator (EHII) [22] (an indicator of the equivalised total disposable household income at baseline) was constructed for the NINFEA participants.

Children's birth weight and gestational age at birth were registered at birth. Weight at 6 months of age were ascertained from the 6-month questionnaire, and weight at 18 months of age were obtained from the 18-month questionnaire. From the corresponding follow-up

questionnaires at 4, 7, and 10 years of age, our NINFEA dataset includes 4232, 2152 and 973 measurements of children's weight and height, respectively. These two measurements were used to calculate each children's body mass index (BMI) at each follow-up. Body mass index (BMI) is computed as the ratio between weight (in kg) and squared height (in meters). Overweight and Obesity for each age were defined according to the official International Obesity Task Force (IOTF) cut-offs [25].

M1: EUROMOD and the Fiscal Reform Scenarios

The first module (M1 – the tax-benefit microsimulation model) uses EUROMOD capabilities to simulate the prospective effects of eight benefit policies on household disposable income. EUROMOD produces an output dataset that contains a population that is the same as the Italian EU-SILC 2010 sample (46788 individuals), but with added information on disposable income for each individual, based on the specific, actual or hypothetical, policy system considered. This aforementioned output dataset provided individual disposable income data that was aggregated for each household in order to compute firstly, the equivalised household size, and later on, the equivalised household income. This adjustment was done in compliance with the modified equivalence scale suggested by the Organisation for Economic Cooperation and Development (OECD) [26]. For reasons of comparability between the Italian EU-SILC 2010 and Italian NINFEA 2011 cohort analyses, we excluded families with more than 7 members or families with no children less than 5 years old.

The simulated fiscal interventions are shown in Table 1. Each pair of simulated policies were implemented with two different levels of intensity regarding benefit amounts, but keeping the same rules for eligibility and recipients among them, with the aim to evaluate potential dose-response effects.

Table 1 Simulated tax-benefit scenarios

Benefit scenarios.

| | Baseline (BS) | Basic Inc | come (BI) | Poverty Reduction (PR) | | New-borns Benefit (NB) | | Child Benefit (CB) | |
|-------------------|------------------|-----------------------------|-----------------------------|------------------------------------------------|------------------------------------------------|------------------------------------------------|------------------------------------------------|------------------------------------------------|------------------------------------------------|
| | | BI1 | BI2 | PR1 | PR2 | NB1 | NB2 | CB1 | CB2 |
| Eligibility | - | all | all | households with per capita income < €500 month | households with per capita income < €500 month | households with per capita income < €500 month | households with per capita income < €500 month | households with per capita income < €500 month | households with per capita income < €500 month |
| Benefit Amount | - | € 100 | €100 | €100 | € 100 | € 500 | € 500 | € 500 | € 500 |
| Periodicity | - | yearly | monthly | yearly | monthly | yearly | monthly | yearly | monthly |
| Recipients | - | all household members | all household members | all household members | all household members | every child < 1 year old | every child < 1 year old | every child < 5 years old | every child < 5 years old |

Note: Each simulated fiscal policy has two levels of intensity keeping other features fixed. The same benefit amount is given once a year or once a month.

Baseline scenario (BS) was simulated applying the actual 2018 Italian fiscal system on the Italian EU-SILC 2010 data. Basic Income Scenarios, BI1 and BI2, consist of benefit amounts of €100 per year, or per month, respectively, to all citizens without eligibility requirements. Poverty Reduction scenarios, PR1 and PR2, simulate poverty-relief interventions of €100 per year, or per month, respectively, for each member of a household with an per capita disposable income of less than €500 per month. New-borns Benefit scenarios, NB1 and NB2, simulate more targeted fiscal interventions of €500 per year, or per month, respectively, for each child less than one year old in households with an equivalised disposable income of less than €500 per month. Child Benefit scenarios, CB1 and CB2, simulate the New-borns Benefits scenarios but with the only difference being that the eligibility rule regarding age threshold for recipients is raised from less than one year of age to five years of age. Consequently, these CB1 and CB2 simulated policies reach a larger number of households than the New-borns

These eight simulated policies with two different intensities allowed to compute marginal benefits, which are a normalised measure of the effectiveness of the different policy instruments, and can be used to compare among the prospective health effects of each alternative.

EUROMOD also provides the overall cost for the public budget, and therefore, after comparison with the baseline, the cost of the changes implemented in the counterfactual scenarios, which can be used to calculate the marginal benefit (health outcome gain compared with the policy cost) of the reforms.

M2: The Concatenated Regression Models in the NINFEA birth cohort

The aim of this module is to estimate regression parameters of interest and the corresponding variance-covariance matrix, that are required later in the third module.

The structure of the concatenated regressions models fitted on the NINFEA data is described in the following equations:

2)
$$BW = \alpha_{bw} + \beta_{bw}GA + \delta_{bw}EHII + \Sigma \beta_{bw_s}X_s + \varepsilon$$

204 3)
$$WT_{6m} = \alpha_{wt6} + \theta_{wt6_1}BW + \theta_{wt6_2}GA + \delta_{wt6}EHII + \Sigma \theta_{wt6_s}X_s + \varepsilon$$

205
$$WT_{18m} = \alpha_{wt18} + \theta_{wt18_1}WT_{6m} + \theta_{wt18_2}BW + \delta_{wt18}EHII + \Sigma \theta_{wt18_s}X_s + \varepsilon$$

206 5)
$$BMI_{48m} = \alpha_{b48} + \beta_{b48_1}WT_{18m} + \beta_{b48_2}WT_{6m} + \delta_{b48}EHII + \Sigma \beta_{b48_5}X_s + \varepsilon$$

7)
$$BMI_{120m} = \alpha_{b120} + \beta_{b120_1}BMI_{84m} + \beta_{b120_2}BMI_{48m} + \delta_{b120}EHII + \Sigma \beta_{b120_s}X_s + \varepsilon$$

GA and BW stand for gestational age and child's birth weight, respectively. WT_{6m} and WT_{18m} stand for child's weight at 6 and 18 months after being born, respectively. BMI_{48m}, BMI_{84m}, and BMI_{120m}, stand for child's body mass index (BMI) at 4, 7 and 10 years of age. As described in detail in the paper by Pizzi et al. [22], an indicator of the EU-SILC-based equivalised total disposable household income (the Equivalised Household Income Indicator - EHII) was constructed for the NINFEA participants within the framework of the H2020 LifeCycle project [27]. In brief, the EHII was constructed using external data provided by the Italian EU-SILC 2011 survey, and individual and household characteristics available in the NINFEA cohort, namely parental age, cohabitation status, education, country of birth and occupation, house size and type, and family size. The EHII is the log transformation of the equivalised household disposable income as used in Pizzi et al. [22]. In all equations EHII is the income indicator (logtransformed), with δ being the estimated coefficient of interest for the income indicator. Moreover, α is the intercept - different for each regression -, $\Sigma \beta_s X_s$ is the sum of sex of child, maternal country of birth, and age at delivery. ε is the error component. The underlying assumption is that all outcomes analysed are influenced by the two previous ones and by the other factors cited above. For each independent variable, these models provide estimated effect sizes, confidence intervals, and the corresponding variance-covariance matrix required by module 3 (M3). M3: Integrating outputs from modules 1 and 2 and creating microsimulation scenarios

The last module of MICH model applies effects sizes from M2 to the population obtained from M1. The effects of EHII on health outcomes, used to simulate the impact of fiscal policies was estimated by the δ coefficients of the multivariable regressions described above. From M1

output we select the population of children less than 5 years old, and expand it using the Italian EU-SILC 2011 survey sample weights, obtaining a study population of 30910 children. Using the same set of concatenated multivariable linear regressions shown above, with the outputs from M1 and M2 used as inputs and the estimated alphas, betas and standard errors of the regressions from M2, the integrated model estimates the distribution of gestational age (GA) for the population of children under 1 year old using regression equation 1. The obtained distribution of gestational age (GA) is successively introduced in regression equation 2 together with the same set of demographic and socioeconomic variables, including the equivalised household disposable income from M1. Regressions equations from 3 to 7 use the same principle, creating a flow of outputs used as inputs for the next regression model and allowing us to simulate the final Body Mass Index (BMI) distributions at 18, 48 and 84 months, in a sequential order. For each outcome and each scenario, 1000 simulations were performed using the Monte Carlo sampling method [28]. This allows the main parameter values, in our case the estimated alphas and betas of the regression equations, to vary in each simulation cycle according to their assumed underlying distributions and their variance-covariance matrix. The number of simulations was chosen after verifying that the estimates were stable and further runs were neither modifying our point estimates, corresponding standard errors, nor other aspects of the simulation. Because the intercepts were obtained from the regression models applied to the NINFEA cohort, which is not representative of the Italian population, for M2 we needed to calibrate their values. The calibration was achieved varying the alpha of the regression model 1

described above, in order to obtain the lower sum of squared errors (SSE) in comparison with

the Italian national prevalence of premature births, and for regression model 2 with the average birth weight, both from the "Certificato di Assistenza al Parto" (CEDAP, translated as Italian birth registry) of the year 2011 [29].

All scenarios were compared in terms of prevalence ratios, using the selected scenario as numerator and the real fiscal scenario as denominator. Marginal benefits were obtained by dividing the cost of the fiscal intervention, provided by EUROMOD in M1, by the prevalence difference between scenarios.

M1 was executed in EUROMOD version 3.0.0, and its output processed in STATA version 14.

M2 and M3 were coded and implemented in R version 3.6.3.

RESULTS

Table 2 shows baseline values corresponding to health outcomes, demographic and socioeconomic variables, as well as simulated parameters, that were used to build our three-part MICH model.

The mean of log of the household equivalised income at baseline in the Italian EU-SILC 2010 population is lower than in the NINFEA cohort (7.02 and 7.38, respectively). However, the percentage of mothers not born in Italy in the Italian EU-SILC 2010 dataset is higher than in the NINFEA cohort (16.8% and 4.2%, respectively). Modules 1 and 2 portrait similar values for the remaining demographic and socioeconomic variables.

Table 3 shows estimated coefficients for each regression model in the series of concatenated regressions included in M2.

Table 2 Estimated means, percentages and parameters used in the M1, M2 and M3 modules of the MICH model

| | M1 | M2 | M3 |
|------------------------------------------|-------------------|----------------|----------------------------|
| | EUROMOD (EU-SILC) | NINFEA Cohort | Baseline Simulated Values* |
| Health Outcomes | | | |
| Gestational Age (weeks) | - | 39.5 [1.8] | 39.6 [0.28] |
| Birth Weight (kg) | - | 3,237 [499] | 3,218 [66] |
| Weight at 6 months (kg) | - | 7,539 [942] | 7,774 [241] |
| Weight at 18 months (kg) | - | 11,162 [1,285] | 11,286 [256] |
| BMI at 48 months | - | 15.6 [1.7] | 15.5 [0.2] |
| BMI at 84 months | - | 15.9 [2.1] | 16.0 [0.5] |
| BMI at 120 months | - | 17.3 [2.6] | 17.9 [1.0] |
| Demographic and Socioeconomic Predictors | | | |
| Female gender | 51.60% | 49.30% | - |
| Log of Equivalised income | 7.02 [0.67] | 7.38 [0.26] | - |
| Foreign citizenship of the mother | 16.80% | 4.20% | - |
| Age of the mother | 33.6 [5.2] | 33.3 [4.4] | - |

Note: Weight in kg. Estimated mean values (with standard deviations in brackets), or percentages. *Distribution of the means of the total runs.

Table 3 Estimated coefficients of the concatenated multivariable regressions from Module 2 (M2)

| | (1) Gestational age | (2) Birth | (3) Weight | (4) Weight | (5) BMI | (6) BMI | (7) BMI |
|----------------------------------|--------------------------|-------------------------|--------------------------|--------------------------|-------------------------|-------------------------|-------------------------|
| VARIABLES | at birth (weeks) | weight (kg) | at 6 months (kg) | at 18 months (kg) | at 48 months | at 84 months | at 120 months |
| EHII | 0.31 | -0.55 | -0.60 | 1.36 | -0.31 | -0.52 | -0.85 |
| Mother's age | [0.11 - 0.51] -0.05 | [-0.990.10] 0.03 | [-1.58 - 0.39] -0.05 | [0.07 - 2.65] -0.02 | [-0.570.05] -0.00 | [-0.930.10] 0.00 | [-1.580.12] -0.01 |
| · · | [-0.060.04] | [0.01 - 0.06] | [-0.110.00] | [-0.09 - 0.05] | [-0.02 - 0.01] | [-0.02 - 0.03] | [-0.05 - 0.03] |
| Mother's country of birth | 0.02 [-0.19 - 0.22] | 0.65 [0.11 - 1.19] | 1.88 [0.66 - 3.09] | -1.30 [-2.82 - 0.23] | -0.20 [-0.54 - 0.14] | -0.01 [-0.62 - 0.61] | 0.17 [-0.87 - 1.22] |
| Sex | -0.05 [-0.13 - 0.04] | -1.39 [-1.581.19] | -4.46 [-4.904.02] | -1.47 [-2.070.86] | 0.29 [0.18 - 0.41] | 0.16* [-0.01 - 0.34] | -0.12 [-0.40 - 0.17] |
| Gestational age at birth (weeks) | | 1.70 [1.63 - 1.76] | -0.46 [-0.620.29] | | | | |
| Birth weight (kg) | | | 0.89 [0.83 - 0.94] | 0.23 [0.16 - 0.30] | | | |
| Weight at 6 months (kg) | | | | 0.83 [0.79 - 0.87] | 0.01 [0.00 - 0.02] | | |
| Weight at 18 months (kg) | | | | [25 | 0.05 [0.05 - 0.06] | 0.02 [0.01 - 0.03] | |
| Body Mass Index at 48 months | | | | | [0.00 | 0.55 [0.48 - 0.63] | 0.27 [0.15 - 0.39] |
| Body Mass Index at 84 months | | | | | | [0.40 0.03] | 0.77 [0.67 - 0.88] |
| Constant | 38.83 [37.39 - 40.27] | -31.01 [-34.9827.03] | 73.19 [64.52 - 81.86] | 32.69 [22.92 - 42.47] | 10.70 [8.75 - 12.66] | 8.73 [5.55 - 11.92] | 7.62 [2.03 - 13.22] |
| Observations | 6,387 | 6,202 | 5,173 | 4,141 | 2,923 | 1,621 | 658 |
| R-squared | 0.01 | 0.37 | 0.28 | 0.46 | 0.20 | 0.26 | 0.49 |

Note: 95% Confidence Intervals in brackets. EHII stands for Equivalised Household Income Indicator. Sex (0=Male; 1 Female). Mother's country of birth (0=Italy; 1= others)

The significant reduction in terms of number of observations over ages was not due to cohort attrition, but to the dynamic recruitment of the children from the year 2004 onwards, as result most of children recruited in more recent years have not attained the oldest ages, as explained in the Additional file 3.

Table 4 shows prevalence ratios of overweight and obesity between the baseline scenario and the eight combinations of fiscal interventions according to Table 1, for children of 48, 84 and 120 months of age. Reductions in terms of population prevalence for the results of Table 4 are reported in the Additional file 1: Table S3.

Table 4 Prevalence Ratios and prediction intervals for children overweight and obesity at 48, 84 and 120 months

| 30 | | Basic Income | | Poverty Reduction | | New-bor | New-borns Benefit | | Child Benefit | |
|----------|---------------------|---------------|---------------|-------------------|---------------|---------------|-------------------|---------------|---------------|--|
| 31 | | BI1 | BI2 | PR1 | PR2 | NB1 | NB2 | CB1 | CB2 | |
| 32 | 48 months: | | | | | | | | | |
| 33 34 | Children | 0.994 | 0.946 | 0.996 | 0.967 | 0.999 | 0.993 | 0.993 | 0.952 | |
| 35 | Overweight | [0.987-1.000] | [0.895-0.998] | [0.991-1.000] | [0.934-1.001] | [0.998-1.000] | [0.985-1.000] | [0.985-1.001] | [0.904-1.000] | |
| 36 | | | | | | | | | | |
| 37 | Children | 0.986 | 0.907 | 0.988 | 0.936 | 0.998 | 0.987 | 0.982 | 0.912 | |
| 38 | Obesity | [0.966-1.005] | [0.809-1.004] | [0.971-1.006] | [0.861-1.012] | [0.994-1.002] | [0.971-1.004] | [0.957-1.008] | [0.813-1.011] | |
| 39 | · | | . , | . , | , | . , | | | | |
| 40 41 | 84 months: | | | | | | | | | |
| 42 | Children | 0.991 | 0.913 | 0.994 | 0.946 | 0.999 | 0.988 | 0.989 | 0.921 | |
| 43 | Overweight | [0.985-0.996] | [0.858-0.968] | [0.99-0.998] | [0.911-0.982] | [0.997-1.000] | [0.979-0.996] | [0.982-0.996] | [0.869-0.974] | |
| 44 | 5 | | | | , | | | | | |
| 45 | | 0.968 | 0.807 | 0.972 | 0.854 | 0.996 | 0.973 | 0.957 | 0.805 | |
| 46 | Children Obesity | [0.94-0.995] | [0.67-0.945] | [0.947-0.997] | [0.734-0.974] | 0.991-1.002] | [0.949-0.997] | [0.918-0.995] | [0.654-0.957] | |
| 47 48 | Obesity | [0.94-0.993] | [0.07-0.945] | [0.947-0.997] | [0.734-0.974] | 0.991-1.002] | [0.949-0.997] | [0.916-0.993] | [0.034-0.937] | |
| 49 | 120 months: | | | | | | | | | |
| 50 | | 0.988 | 0.876 | 0.992 | 0.925 | 0.998 | 0.982 | 0.986 | 0.887 | |
| 51 | Children | | | | | | | | | |
| 52 | Overweight | [0.982-0.993] | [0.819-0.933] | [0.988-0.996] | [0.888-0.961] | [0.997-0.999] | [0.972-0.992] | [0.98-0.993] | [0.831-0.943] | |
| 53 | | | | | | | | | | |
| 54 | Children | 0.946 | 0.666 | 0.951 | 0.721 | 0.993 | 0.950 | 0.920 | 0.639 | |
| 55 56 | Obesity | [0.918-0.975] | [0.501-0.83] | [0.924-0.979] | [0.562-0.881] | [0.985-1.001] | [0.917-0.982] | [0.875-0.965] | [0.439-0.838] | |

57 Note: 95% Confidence intervals in brackets. Ratios and prediction intervals according to the different fiscal reform scenarios in 58 comparison with the baseline.

Firstly, we observe that Child benefit intervention CB2, consisting of a monthly benefit amount of €500 for each child younger than 5 years in families with household equivalised disposable income lower than €500, has very strong effects on overweight (-11%) and obesity (-36%) for children of 120 months of age, with a prevalence rate of 0.89 (95%CI: 0.83-0.94) and 0.64 (95%CI: 0.44-0.84), respectively. Secondly, Basic Income intervention BI2, consisting of a monthly benefit amount of €100 for each family member and no further eligibility requirements, again for children of 120 months of age, shows quite important effects on overweight (-12%) and obesity (-33%) with a prevalence rate of 0.88 (95%CI: 0.82-0.93) for the former outcome variable of interest, and 0.67 (95%CI: 0.50-0.83) for the latter.

Figure 2 shows the distribution of the equivalised disposable income in logarithmic units, and the distribution of Body Mass Index (BMI) at different ages. First column of figures portraits the less focalized intervention, Basic Income BI2, whereas second column depicts the more focalized intervention, Child Benefit CB2. Black curve represents distribution at baseline scenario, whereas the red curve represents the distribution after the benefit policy simulation.

Basic Income BI2 simulation shows a similar pattern to Child Benefit CB2 simulation, although shifts for the latter seem to be more distinct than those for the former. Child Benefit CB2 simulation shows a correspondence between the shift to the right on the equivalised income distribution and the subsequent shift to the left on BMI distributions at all ages.

Finally, Table 5 shows estimated costs for 1% prevalence reduction in overweight and obesity at 48, 84 and 120 months of age with respect to the baseline scenario. It's noteworthy to mention that another salient feature of EUROMOD is that it automatically computes and

provides the prospective cost of each simulated scenario. Given the different characteristics of the simulated benefit policies, our estimated costs computations for 1% prevalence reduction show a wide range of values. At 48 months after birth, NB1 and CB1 policies are the most efficient scenarios for both, overweight and obesity, with estimated costs of €6.4 and €27.6 billion, respectively, for NB1, and €4.7 and €15.0 billion, respectively, for CB1. At 84 months after birth the most favourable policies are the same as those at 48 months after birth. NB1 shows costs of €2.4 and €10.7 billion for overweight and obesity, respectively, whereas CB1 depicts costs of €2.0 and €5.5 billion for the aforementioned variables, respectively. Finally, at 120 months after birth, again, NB1 and CB1 scenarios are the most efficient in order to reduce prevalence in 1% with respect to baseline scenario. For overweigh and obesity, NB1 shows costs of €1.4 and €4.6 billion, respectively, whereas CB1 scenario shows costs of €1.2 and €2.7 billion, respectively. It is worthy to point out that, according to results shown in Table 4 and Table 5, we observe

that at 48, 84, and 120 months after birth, cost estimates for NB1 and NB2 are distinctly smaller than those corresponding to Basic Income scenarios BI1 and BI2. However, estimated reductions in prevalence ratios among BI1 and New-borns Benefit scenarios, NB1 and NB2, are rather similar.

56 341

Figure 2 Kernel Density plots of the distribution of the logarithm of equivalised disposable

income, and BMI at 48, 84, and 120 months of age

BI2 CB₂ 0.8 0.8 9.0 0.4 (Log) 0.4 0.2 0.2 0.0 2.5 2.0 2.0 BMI 48m 1.0 9.0 1.2 0. 0.8 BMI 84m 0.5 0.2 0.0 9.0 9.0 0.5 0.4 BMI 120m 0.2 0.2 0.1

Note: Black curve = distribution at baseline. Red curve= distribution after corresponding benefit policy simulation. Bl2= Simulated Fiscal Policy of Basic Income for all families with a cash transfer of epsilon 100 per month to each family member. CB2= Simulated Fiscal Policy of Child Benefit given to households with per capita income epsilon 500 month, being the cash transfer of epsilon500 monthly for each child epsilon5 years old.

Table 5 Estimated costs for 1% prevalence reduction in overweight and obesity

| | Basic Income | | Poverty | Poverty Reduction | | New-borns Benefit | | Child Benefit | |
|-------------|--------------|-------|---------|-------------------|------|-------------------|------|---------------|--|
| | BI1 | BI2 | PR1 | PR2 | NB1 | NB2 | CB1 | CB2 | |
| 48 months: | | | | | | | | | |
| Overweight | 79.8 | 109.9 | 20.6 | 32.5 | 6.4 | 8.7 | 4.7 | 8.4 | |
| Obesity | 272.1 | 505.0 | 60.7 | 133.2 | 27.6 | 40.5 | 15.0 | 36.4 | |
| 84 months: | | | | | | | | | |
| Overweight | 33.7 | 43.1 | 9.0 | 12.7 | 2.4 | 3.3 | 2.0 | 3.3 | |
| Obesity | 109.0 | 217.1 | 22.7 | 51.5 | 10.7 | 17.0 | 5.5 | 14.6 | |
| 120 months: | | | | | | | | | |
| Overweight | 19.7 | 23.3 | 5.6 | 7.0 | 1.4 | 1.8 | 1.2 | 1.8 | |
| Obesity | 59.3 | 110.2 | 11.9 | 23.8 | 4.6 | 7.9 | 2.7 | 6.9 | |

Note: Prevalence reduction with respect to baseline scenario. Figures in billions of euros.

DISCUSSION

This study shows how simulated fiscal reforms, and in particular poverty-reduction fiscal policies, could strongly reduce overweight and childhood obesity in a high-income European country, such as Italy. Our findings quantify also the dose-response relationship between increased benefits and impact on overweight and obesity for each of the eight simulated fiscal interventions. Moreover, focalising the simulated interventions on households with newborns or children, instead of on all households, seemed to be particularly efficient in terms of marginal benefits.

To our knowledge this is the first study that creates a comprehensive microsimulation model to evaluate the effects of fiscal policies on health, taking advantage of a consolidated platform such as EUROMOD, and integrating it with microsimulation algorithms that project the effects of the equivalised disposable household income on the chosen health outcome. Ultimately, our approach allows to forecast the effectiveness and efficiency of large fiscal interventions on a representative sample of the population.

 Besides its methodological sophistication and comprehensiveness, the MICH model constitutes an unprecedented attempt to provide evidence of how fiscal policies could affect health outcomes in the population, and could be a systematically used tool in fiscal and health policy-making. Only another recently published study used EUROMOD to evaluate the impact of fiscal policies on overall mortality in Scotland, showing that policies targeting the poorest populations were the most effective to reduce inequalities [30]. However, while its fiscal simulations were comprehensive, the association between income and mortality - used to develop all forecasting scenarios - was not obtained through individual-level longitudinal data, but it was estimated by a cross-sectional regression model based only on population quintiles.

Fiscal reforms have historically been evaluated only in terms of their economic effects, while their impact on other dimensions of society have been neglected [31]. The MICH model could provide evidence of such effects at the time of decision-making, allowing a more balanced and informed choice and implementation of fiscal policies. Moreover, the model is particularly relevant in high-income countries where poverty-alleviation policies are mainly based on fiscal reforms and are often required to demonstrate effective targeting and efficiency of fiscal benefits to the poorest populations. Finally, being based on data from the EUROMOD platform and from the EU-SILC surveys, which are available for several European countries, and being the EHII available for several European birth cohort studies (almost all LifeCycle cohorts [27]), the MICH model could be used to evaluate and compare the effect of a broad range of fiscal policies on childhood health across different European countries.

The choice of the fiscal policies in the simulation has been based on the most common universal and targeted poverty-reduction interventions currently implemented by the

majority of EU governments. Household disposable income is among the strongest social determinant of health, because it is a direct measure of material resources, and changes in its levels could have an effect on several health outcomes for the members of the family [2, 32–34]. Previous studies have shown an association between household income and child obesity [35, 36], suggesting the implementation of income subsidies to reduce the economic restrictions of those individuals with lesser economic means. To the best of our knowledge there are no randomized trials of fiscal interventions or increased total family disposable income with child BMI as the outcome, but a few quasi-experimental studies have evaluated the impact of cash transfers on child obesity. [37] The overall impact of these interventions seems to be dependent on the baseline income of the intervention recipients, the age of children at the time of transfer and also the context where the intervention is delivered [38–40].

Other mathematical modelling studies have attempted to evaluate the effects of different interventions on childhood obesity. A recent microsimulation study showed the combined effectiveness of after-school physical activities, sugar-sweetened beverage taxes, and a ban on child-directed fast-food advertising [8]; another study created a specific microsimulation model called Early Prevention of Obesity in Childhood (EPOCH) which was able to model BMI trajectories from early childhood to adolescence [10]. Several other simulation models included behavioural and environmental contributors, and focused policy interventions [9]. However, none of these foregoing studies included income, and its variations due to poverty-reduction interventions, as one of the explanatory variables of childhood obesity.

This study and its microsimulation model are particularly relevant during the current COVID-19 pandemic, which has triggered the worst global economic crisis since the Great

Depression, with a World GDP contraction of 4.5% and dramatic increases in unemployment and poverty rates in almost all nations [41, 42].

One of the main instruments used by most governments to increase the resilience of their populations has been extended fiscal benefits, but is still unclear how much these interventions have been able to mitigate the increase in household poverty levels. In fact, the strict or partial lockdowns applied in almost all countries could have increased sedentary behaviours, and the dramatic income losses could have forced some families to buy cheaper and more obesogenic foods, especially for children. While physical activity interventions would be difficult and slow to implement due to lockdown measures, fiscal poverty-reduction interventions could be a fast and effective instrument to curb this tendency.

One of the main limitations of this study regards the estimated marginal effects. Fiscal reforms have a wide range of effects, spanning from socioeconomic to health outcomes, and the objective of our study is to show their - mostly unintended - impact only on a specific health outcome, child overweight and obesity. As a consequence, the exercise of evaluating the marginal benefits cannot be interpreted as a cost-effectiveness evaluation. Another limitation of our study is that, while our statistical associations, obtained from a cohort study, are respecting many of the traditional conditions of causality (strength of the associations, consistency, specificity, temporality and plausibility), and have the advantage of being specific for the population in which the simulations were conducted, the observational nature of the study cannot rule out the possibility that they are not causal. We also acknowledge that the NINFEA cohort has attrition, even if within the standard magnitude for such kind of child cohorts. Moreover, we did not include all the variables associated with overweight and obesity in the concatenated regression models of MICH M2 module. This is because we

included confounding factors only of the association between EHII and BMI, and not mediators of such association (such as childhood diet, physical activity, etc.) that would have affected the real effect estimates of EHII. However, due to the complexity of the overweight and obesity theoretical framework, we acknowledge that we cannot completely rule out the possibility of remaining omitted variable bias.

CONCLUSIONS

In conclusion, our study illustrates the construction of an unprecedented integrated microsimulation model — based on the consolidated EUROMOD platform - able to forecast the effect of a broad range of fiscal interventions on childhood obesity, with algorithms and codes potentially flexible to be used for other health outcomes and for other European countries. While the impact of fiscal policies is usually measured on economic outcomes, our study was the first to quantify their effects on one of the most concerning child health problems in high-income countries. Potential impacts of fiscal interventions on health of the population should be taken into account during the process of policy-making, and should be considered in the framework of the Health in All policies (HiAP) [11].

| | 452 | LIST OF ABREVIATIONS |
|----------|-----|---------------------------------------------------------------------------------------|
| 1 2 | 453 | BI: Universal Basic income |
| 3 4 | 454 | BI1: Baseline Income scenario 1 |
| 5 6 | 455 | BI2: Baseline Income scenario 2 |
| 7 8 | 456 | BMI: Body Mass Index |
| 9 10 | 457 | BMI _{48m} : Child's body mass index at 4 years of age |
| 11 12 | 458 | BMI _{84m} : Child's body mass index at 7 years of age |
| | 459 | BMI _{120m} : Child's body mass index at 10 years of age |
| 15 16 | 460 | BS: Baseline Scenario |
| 17 18 | 461 | BW: Child's birth weight |
| 19 20 | 462 | CB: Child Benefit |
| 21 22 | 463 | CB1: Child Benefit scenario 1 |
| 23 24 | 464 | CB2: Child Benefit scenario 2 |
| 25 26 | 465 | CEDAP: Certificato di Assistenza al Parto (translated as Italian birth registry) |
| 27 28 | 466 | COVID-19: Coronavirus Disease 2019 |
| 29 30 | 467 | DG EMPL: Directorate-General for Employment, Social Affairs and Inclusion |
| 31 32 | 468 | EHII: Equivalised Household Income Indicator |
| | 469 | EPOCH: Early Prevention of Obesity in Childhood |
| 35 36 | 470 | EU: European Union |
| 37 38 | 471 | EU-SILC: European Union Statistics on Income and Living Conditions |
| 39 40 | 472 | GA: Gestational Age |
| 41 42 | 473 | GDP: Gross Domestic Product |
| 43 44 | 474 | H2020: Horizon 2020 |
| | 475 | HiAP: Health in All Policies |
| 47 48 | 476 | IOTF: International Obesity Task Force |
| 49 50 | 477 | ISPOR-SMDM: International Society for Pharmacoeconomics and Outcomes Research and the |
| 51 52 | 478 | Society for Medical Decision Making |
| 53 54 | 479 | MICH: Microsimulation for Income and Child Health |
| 55 56 | 480 | M1: Module 1 |
| 57 58 | 481 | M2: Module 2 |
| 59 60 | 482 | M3: Module 3 |
| 61 62 | | |
| 63 64 | | |
| 65 | | 25 |

NINFEA: Nascita e Infanzia: gli Effetti dell'Ambiente (translated as Birth and Childhood: The Effects of the Environment)

NB1: New-borns Benefit scenario 1

NB2: New-borns Benefit scenario 2

OECD: Organisation for Economic Cooperation and Development

488 PR: Prevalence Ratio

489 PR1: Poverty Reduction scenario 1

490 PR2: Poverty Reduction scenario 2

SEP: Socio-Economic Position

UK: United Kingdom

493 WT_{6m}: Child's weight at 6 months after being born

494 WT_{18m}: Child's weight at 18 months after being born

DECLARATIONS

Acknowledgments

The authors are grateful to all participants of the NINFEA cohort. EUROMOD is maintained, developed and managed jointly by the Institute for Social and Economic Research (ISER) at the University of Essex and the Joint Research Centre of the European Commission, in collaboration with national teams from the EU member states. We are indebted to the many people who have contributed to the development of EUROMOD. The process of extending and updating EUROMOD is financially supported by the European Union Programme for Employment and Social Innovation 'Easi' (2014-2020). The results and their interpretation are the authors' responsibility. We are grateful for the valuable comments of Prof. Franco Sassi. DR and NB acknowledge support from the Spanish Ministry of Science and Innovation and State Research Agency through the "Centro de Excelencia Severo Ochoa 2019-2023" Program (CEX2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program.

Consent for Publication

Not applicable.

Funding

LIFECYCLE. The LIFE-CYCLE project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 733206. This publication reflects only the author's views and the European Commission is not liable for any use that may be made of the information contained therein. STOP. The STOP project (http://www.stopchildobesity.eu/) received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No.774548. The STOP Consortium is coordinated by Imperial College London and includes 24 organisations across Europe, the United States and New Zealand. The content of this publication reflects only the views of the authors, and the European Commission is not liable for any use that may be made of the information it contains. NINFEA. The NINFEA cohort was partially funded by the Compagnia San Paolo Fundation and by the Piedmont Region. MAH is supported by the British **Heart Foundation.**

Availability of Data and Materials

Anonymus individual data from the NINFEA Cohort could be provided upon request to the authors. EU-SILC dataset used as input for EUROMOD are available upon request to Eurostat (https://ec.europa.eu/eurostat) and to EUROMOD (https://euromod-web.jrc.ec.europa.eu/)

REFERENCES

2 3

1

549

548

- 5
- б
- 9 10 11
- 13 ¹⁴ 555
- 15 16
- 18
- ¹⁹ 558 20
- 21 22
- 23
- 24 561 25
- 26 27
- ₂₈ 563
- 30 31
- 32 33
- 34 **567** 35 36
- 37 38
- ⁴⁰ 571 41 42
- 43 44 573
- 46 47 48
- 49 576 51
- 53 55
- 57 58 59
- 56
- 60 61 62

- 1. Phipps SA, Burton PS, Osberg LS, Lethbridge LN. Poverty and the extent of child obesity in
- 550 Canada, Norway and the United States. Obes Rev Off J Int Assoc Study Obes. 2006;7:5–12.
- 2. Marmot M, Friel S, Bell R, Houweling TA, Taylor S. Closing the gap in a generation: health 7 551
 - 552 equity through action on the social determinants of health. The Lancet. 2008;372:1661–9.
 - 553 3. Lenhart O. The effects of income on health: new evidence from the Earned Income Tax
- 12 554 Credit. Rev Econ Househ. 2019;17:377-410.
 - 4. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of
 - 556 1698 population-based measurement studies with 19.2 million participants. The Lancet.
- 17 557 2016;387:1377-96.
 - 5. Greves Grow HM, Cook AJ, Arterburn DE, Saelens BE, Drewnowski A, Lozano P. Child
 - 559 obesity associated with social disadvantage of children's neighborhoods. Soc Sci Med.
 - 560 2010;71:584–91. doi:10.1016/j.socscimed.2010.04.018.
 - Gable S, Lutz S. Household, Parent, and Child Contributions to Childhood Obesity. Fam
 - 562 Relat. 2000;49:293–300. doi:https://doi.org/10.1111/j.1741-3729.2000.00293.x.
 - 7. Ogden CL, Carroll MD, Fakhouri TH, Hales CM, Fryar CD, Li X, et al. Prevalence of Obesity
- 29 564 Among Youths by Household Income and Education Level of Head of Household — United
 - 565 States 2011–2014. Morb Mortal Wkly Rep. 2018;67:186–9.
 - 566 doi:10.15585/mmwr.mm6706a3.
 - 8. Kristensen AH, Flottemesch TJ, Maciosek MV, Jenson J, Barclay G, Ashe M, et al. Reducing
 - 568 childhood obesity through U.S. federal policy: a microsimulation analysis. Am J Prev Med.
 - 569 2014;47:604-12.
- 39 **570** 9. Levy DT, Mabry PL, Wang YC, Gortmaker S, Huang TT-K, Marsh T, et al. Simulation models
 - of obesity: a review of the literature and implications for research and policy. Obes Rev.
 - 572 2011;12:378-94.
 - 10. Hayes A, Tan EJ, Lung T, Brown V, Moodie M, Baur L. A New Model for Evaluation of
- 45 574 Interventions to Prevent Obesity in Early Childhood. Front Endocrinol. 2019;10.
 - 575 doi:10.3389/fendo.2019.00132.
 - 11. World Health Organization, Finland, Sosiaali- ja terveysministeriö, Global Conference on
- 50 **577** Health Promotion. Health in all policies: Helsinki statement, framework for country action:
 - 578 the 8th Global Conference on Health Promotion jointly organized by. 2014.
- 52 http://apps.who.int/iris/bitstream/10665/112636/1/9789241506908 eng.pdf. Accessed 15 579
- 54 580 Apr 2021.
 - 581 12. Bargain O, Orsini K, Peichl A. Comparing Labor Supply Elasticities in Europe and the 582
 - United States New Results. J Hum Resour. 2014;49:723–838. doi:10.3368/jhr.49.3.723.

- 13. Barrios S, Dolls M, Maftei A, Peichl A, Riscado S, Varga J, et al. Dynamic Scoring of Tax
- Reforms in the European Union. J Policy Anal Manage. 2019;38:239-62.
 - doi:https://doi.org/10.1002/pam.22105.

- **586** 14. The design of fiscal consolidation measures in the European Union: distributional effects
- and implications for macroeconomic recovery - Institute for Social and Economic Research
 - (ISER). https://www.iser.essex.ac.uk/research/publications/working-papers/euromod/em5-
- 15. Accessed 22 Apr 2021.

- 11 590 15. Decoster A, Jukka Pirttilä J, Holly Sutherland H, Gemma Wright G. SOUTHMOD:
- Modelling tax-benefit systems in developing countries. Int J Microsimulation. 2019;12:1–12.

- 15 592 16. Caro JJ, Briggs AH, Siebert U, Kuntz KM, Force I-SMGRPT. Modeling good research
- 16 593 practices—overview: a report of the ISPOR-SMDM Modeling Good Research Practices Task
- Force-1. Med Decis Mak Int J Soc Med Decis Mak. 2012;32:667–77.

- 17. Sutherland H, Figari F. EUROMOD: the European Union tax-benefit microsimulation 20 595
- 21 596 model. Int J Microsimulation. 2012;6:4–26. doi:10.34196/ijm.00075.

- ²³ 597 18. Eurostat. U - Statistics on Income and Living Conditions microdata 2004-2018, release
- 2020, version 1. 2020.
- https://ec.europa.eu/eurostat/documents/203647/203704/EU+SILC+DOI+2020v1.pdf. 26 599
- Accessed 15 Apr 2021.

- 19. Ceriani L, Figari F, Fiorio C. EUROMOD Country Report ITALY (IT) 2017-2020. 2020.
- 31 602 https://euromod-web.jrc.ec.europa.eu/using-euromod/country-reports/latest. Accessed 15
- ³² 603 Apr 2021.

- 20. Deza M, Jara HX, Oliva N, Torres J. Simulating Personal Income Tax Reforms and Fiscal
- Gains in the Andean Region. Inter-American Development Bank; 2020. 36 605
- doi:10.18235/0002493.

- 21. EUROMOD. Centre for Microsimulation and Policy Analysis.
- https://www.microsimulation.ac.uk/euromod/. Accessed 20 Sep 2021.

- 43 609 22. Pizzi C, Richiardi M, Charles M-A, Heude B, Lanoe J-L, Lioret S, et al. Measuring Child
- Socio-Economic Position in Birth Cohort Research: The Development of a Novel
- Standardized Household Income Indicator. Int J Environ Res Public Health. 2020;17:1700.
- 48 612
 - 23. Richiardi L, Baussano I, Vizzini L, Douwes J, Pearce N, Merletti F, et al. Feasibility of recruiting a birth cohort through the Internet: the experience of the NINFEA cohort. Eur J
 - Epidemiol. 2007;22:831-7.

24. Progetto NINFEA | Index. https://www.progettoninfea.it/. Accessed 22 Apr 2021. **615**

- 25. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness,
- overweight and obesity. Pediatr Obes. 2012;7:284–94.

27. LifeCycle. LifeCycle. https://lifecycle-project.eu/. Accessed 15 Apr 2021.

28. Mooney CZ. Monte Carlo Simulation. SAGE; 1997.

29. Salute della M. Certificato di assistenza al parto (CeDAP). Analisi dell'evento nascita -

Anno 2017.

http://www.salute.gov.it/portale/documentazione/p6 2 2 1.jsp?lingua=italiano&id=2931. 11 624

¹² **625**

Accessed 16 Nov 2020.

- 30. Richardson E, Fenton L, Parkinson J, Pulford A, Taulbut M, McCartney G, et al. The effect
- of income-based policies on mortality inequalities in Scotland: a modelling study. Lancet **627**
- Public Health. 2020;5:e150-6.

- 31. Fair Society Healthy Lives (The Marmot Review). Institute of Health Equity.
- 21 630 http://www.instituteofhealthequity.org/resources-reports/fair-society-healthy-lives-the-
- **631** marmot-review. Accessed 15 Apr 2021.

- 32. Stronks K, van de Mheen H, van den Bos J, Mackenbach JP. The interrelationship
- between income, health and employment status. Int J Epidemiol. 1997;26:592–600.

634

33. Mackenbach JP, Looman CW. Life expectancy and national income in Europe, 1900-

2008: an update of Preston's analysis. Int J Epidemiol. 2013;42:1100–10.

- 34. Cole BL, Shimkhada R, Morgenstern H, Kominski G, Fielding JE, Wu S. Projected health
- **637** impact of the Los Angeles City living wage ordinance. J Epidemiol Community Health.
- 2005;59:645-50.

35. Costa-Font J, Gil J. Intergenerational and socioeconomic gradients of child obesity. Soc Sci Med. 2013;93:29-37.

- 36. Eagle TF, Sheetz A, Gurm R, Woodward AC, Kline-Rogers E, Leibowitz R, et al.
- Understanding childhood obesity in America: Linkages between household income,
- community resources, and children's behaviors. Am Heart J. 2012;163:836-43.

37. Cooper K, Stewart K. Does Money Affect Children's Outcomes? An update. :41.

38. Watson B, Guettabi M, Reimer M. Universal Cash Transfers Reduce Childhood Obesity

49 646

Rates. SSRN Scholarly Paper. Rochester, NY: Social Science Research Network; 2019. doi:10.2139/ssrn.3380033.

39. Akee R, Simeonova E, Copeland W, Angold A, Costello EJ. Young Adult Obesity and

54 649 Household Income: Effects of Unconditional Cash Transfers. Am Econ J Appl Econ. 2013;5:1-

- 40. Akee R, Simeonova E, Copeland W, Angold A, Costello J. Does More Money Make You
- Fat? The Effects of Quasi-Experimental Income Transfers on Adolescent and Young Adult

| 1 2 | 653 654 | Obesity. SSRN Scholarly Paper. Rochester, NY: Social Science Research Network; 2010. https://papers.ssrn.com/abstract=1663175. Accessed 29 Sep 2021. |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| 3 4 5 6 7 8 9 | 655 656 | 41. Han J, Meyer BD, Sullivan JX. Income and Poverty in the COVID-19 Pandemic. National Bureau of Economic Research; 2020. doi:10.3386/w27729. |
| | 657 658 | 42. OECD Economic Outlook, Interim Report September 2020. 2020. http://www.oecd.org/economic-outlook/september-2020. |
| 10 11 12 13 14 15 16 17 18 19 20 12 22 23 24 24 25 26 27 28 29 30 31 31 33 33 34 44 44 44 44 44 44 44 44 44 44 | 659 | |

BMED-D-21-01486R1

Developing an integrated microsimulation model for the impact of fiscal policies on child health in Europe: the example of childhood obesity in Italy Davide Rasella; Lorenzo Richiardi; Nicolai Brachowicz; H. Xavier Jara; Mark Hanson; Delia Boccia; Matteo G. Richiardi; Costanza Pizzi BMC Medicine

Dear Prof. Rasella,

Your manuscript 'Developing an integrated microsimulation model for the impact of fiscal policies on child health in Europe: the example of childhood obesity in Italy' (BMED-D-21-01486R1) has been assessed by our reviewers. Based on these reports, and my own assessment as Editor, I am pleased to inform you that it is potentially acceptable for publication in BMC Medicine, once you have carried out some essential revisions suggested by our reviewers.

Their reports, together with any other comments, are below. Please also take a moment to check our website at

https://www.editorialmanager.com/bmed/l.asp?i=492207&l=8NUGQF0T for any additional comments that were saved as attachments. Once you have made the necessary corrections, please submit a revised manuscript online at:

https://www.editorialmanager.com/bmed/

If you have forgotten your username or password please use the 'Send Login Details' link to get your login information. For security reasons, your password will be reset.

We request that a point-by-point response letter accompanies your revised manuscript. This letter must provide a detailed response to each reviewer/editorial point raised, describing what amendments have been made to the manuscript text and where these can be found (e.g. Methods section, line 12, page 5). If you disagree with any comments raised, please provide a detailed rebuttal to help explain and justify your decision.

Please also ensure that your revised manuscript conforms to the journal style, which can be found in the Instructions for Authors on the journal homepage.

A decision will be made once we have received your revised manuscript, which we expect by 28 Sep 2021.

Please note that you will not be able to add, remove, or change the order of authors once the editor has accepted your manuscript for publication.

Any proposed changes to the authorship must be requested during peer-review, and adhere to our criteria for authorship as outlined in BioMed Central's policies.

To request a change in authorship, please download the 'Request for change in authorship form' which can be found here

- http://www.biomedcentral.com/about/editorialpolicies#authorship.

Please note that incomplete forms will be rejected.

Your request will be taken into consideration by the editor, and you will be advised whether any changes will be permitted.

Please be aware that we may investigate, or ask your institute to investigate, any

unauthorized attempts to change authorship or discrepancies in authorship between the submitted and revised versions of your manuscript.

Once you have completed and returned the form, your request will be considered and you will be advised whether the requested changes will be allowed.

By resubmitting your manuscript you confirm that all author details on the revised version are correct, that all authors have agreed to authorship and order of authorship for this manuscript and that all authors have the appropriate permissions and rights to the reported data.

We look forward to receiving your revised manuscript and please do not hesitate to contact us if you have any questions.

Best wishes,

Ming Yang, PhD BMC Medicine https://bmcmedicine.biomedcentral.com/

Editor's Comments:

1. We require confirmation of authorship from all co-authors, and have not yet received a response from the listed authors. I have resent them the verification email containing the confirmation link. May I kindly ask that you follow up on this or provide us with alternative email addresses, if necessary? Co-authors should contact us directly to confirm co-authorship if they have any problems.

Still to confirm co-authorship

- Delia Boccia
- Matteo G. Richiardi

A: Both co-authors should have confirmed the coauthorship.

2. Please include a figure legend. Figure legends should be no more than 300 words (advised but not mandatory).

A: Now, all figures have the corresponding legend. Section Methods, Figure 1, lines 111-113, page 6, and lines 344-347 page 19.

3. Each file (whether main or Additional) should be mentioned in the main manuscript text in order.

A: Now, all files are mentioned in the main manuscript text in order. Section Methods, lines 104, page 5; line 125, page 6; and line 138, page 7.

4. Please rename supplementary files as 'Additional file X', and cite explicitly by additional file name in the manuscript e.g. 'Additional file 1: Fig. S1'. Please ensure that

if you have more than one additional file that they are cited in ascending order within the main body of text.

- A: Supplementary files were renamed in accordance with BMC Medicine guidelines. Supplementary information Section, lines 530, 532, 534, page 29.
- 5. Please add a description of your additional file to the end of the manuscript. Legends to Combined Additional files should include a short description of the constituent parts, e.g. "Additional File 1: Figures S1-S3. FigS1- [short title]. FigS2 [short title]. FigS3 [short title]...".Please note that legends are advised to be no more than 300 words, though this is not mandatory.
- A: Description of additional files added in Supplementary information Section, lines 531, 533, 535, page 29.
- 6. Please could you update the reference formatting to remove the extra details of Month of publication, date cited, and "Internet". References should follow the format: Smith JJ. The world of science. Am J Sci. 1999;36:234-5. Journal titles should be abbreviated as in the National Library of Medicine https://www.ncbi.nlm.nih.gov/nlmcatalog/journals
- A: Thank you for noticing that, we have changed the references style in accordance with BMC Medicine guidelines.
- 7. Please move all references to the main reference list (i.e. no additional references in Additional Files). These should be included in the main text when the Additional File is first introduced, for example "...see Additional File 1: Table S1) [XX-XX]"; please ensure the reference list is numbered in the order as they appear in the text.
- A: References that were previously included in Supplementary material but not were mentioned in the main manuscript are now cited in Section Methods, lines 104-105, page 5.
- 8. Please make the following change to your section headings: change 'Introduction' to 'Background'.
- A: We did it. Background Section, line 64, page 4.
- 9. Manuscripts should have the following subsections: Background/Methods/Results/Discussion/Conclusions. Please add a new subsection entitled Conclusions.
- A: We added Conclusions subtitle before the already existing conclusion paragraph. Conclusions Section, line 431, page 24.
- 10. Please provide a list of all the abbreviations used in the manuscript. Please place this list just before the Declarations section. If abbreviations are used in the text, they should be defined in the text at first use and included in this list.
- A: List of Abbreviations was added just before the Declarations section, lines 442-484, pages 24-26. For the sake of simplicity and clarity, the Microsimulation for Income and

Child Health Outcomes (MICHO) was renamed Microsimulation for Income and Child Health (MICH) along the text and in the additional files.

- 11. You should include a "Declarations" section, consisting of the following subsections: Acknowledgments; Consent for Publication; Funding; Availability of Data and Materials; Authors' Contributions; Competing interests; Ethics Approval and Consent to Participate. All of these subsections MUST be present.
- The "Authors' Contributions" section of the Declarations should include the text "All authors read and approved the final manuscript"
- The "Competing Interests" section of the Declarations should include any relevant competing interests, or state "The authors declare that they have no competing interests". See here for our editorial policies and a full explanation of competing interests. https://www.biomedcentral.com/getpublished/editorial-policies#CompetingInterests

A: As requested we added the Declarations Section, with all subsections present, from line 489 of page 26.

Reviewer reports:

Reviewer #1: This is a revised manuscript describing a microsimulation model testing the impact of a range of fiscal policies on child weight in Italy. The authors have done a good job of addressing the issues raised by myself and the other reviewer. I just have two remaining comments.

The first relates to a comment in the response to reviewers: "To the best of our knowledge there are no randomized trials of fiscal interventions or increased total family disposable income with child BMI as the outcome. There are several quasi experimental studies on cash transfer, but those studies were conducted in a different population (United States) and did not include child BMI as one of the health outcomes." There are actually a handful of quasi experimental studies on cash transfers with BMI as the outcome, across several countries, and these have found mixed results - see review article: Cooper K, Stewart K. (2017) Does money affect children's outcomes? An update. Case paper 203; ISSN 1460-5023. Centre for Analysis of Social Exclusion, London.

ttp://eprints.lse.ac.uk/103494/1/casepaper203.pdf. This should be acknowledged in light of the strong effects implied by the microsimulation.

A: We thank the reviewer for bringing to our attention the review from Cooper et al. In their work they only mention one study about EITC and obesity from Chia et al, 2013, which admittedly we did not know. As suggested by the reviewer, there are indeed a handful of studies on the effect of income support interventions and obesity among children; however, conclusions from these studies need to be interpreted cautiously, because: a. Only few come from high income countries (the vast majority are from middle income countries); b. results can be conflicting as in some cases these interventions have been associated with an increase of obesity among children. This seems to be dependent on the baseline income of the intervention recipients, the age of children at the time of transfer and also the context where the intervention is delivered). Interestingly, in LMICS results become more consistent in the direction of a positive impact when interventions are delivered in combination with health education and in context where food quality and availability is improved. Overall, these findings

do not undermine the role of income-support interventions in addressing child obesity, but suggest that their maximum potential is achieved when delivered in combination with health education and within a food secure environment (in terms of food quality).

We have added at page 22 line 391: "To the best of our knowledge there are no randomized trials of fiscal interventions or increased total family disposable income with child BMI as the outcome, but a few quasi-experimental studies have evaluated the impact of cash transfers on child obesity. [37] The overall impact of these interventions seems to be dependent on the baseline income of the intervention recipients, the age of children at the time of transfer and also the context where the intervention is delivered [38–40]."

Second, there was another comment in the response to reviewers, "response rates and attrition are at the standard level (after 10 years a lost of around 30% of the participants). Due to that reason, and that the attrition was not differential due to the overall characteristics of the remaining individuals, we didn't use any approach to correct for that." The table in the Supplementary Methods described the level of attrition - thank you - but I couldn't see evidence that the attrition was not differential, though perhaps I missed it. Can this be made clear and a brief comment included in the paper indicating that attrition was not differential?

A: We thank the reviewer for this comment. The sentence about non-differentiality was based on the comparison of the distribution of the weight measures at 18 months between those lost to the 4-year follow-up and those still in the study at that time: mean weight at 18 months in the two groups equal to 11130 and 11191 grams respectively (p-value from T-Student test=0.23). However, we agree with the reviewer that the statement about non-differentiability is too strong as several other factors might affect attrition, so we did not included such statement in the main text or in the additional files, and we have left in the limitations section (p.24 line 433):" We also acknowledge that the NINFEA cohort has attrition, even if within the standard magnitude for such kind of child cohorts."

Editorial Policies

.....

Please read the following information and revise your manuscript as necessary. If your manuscript does not adhere to our editorial requirements, this may cause a delay while this is addressed. Failure to adhere to our policies may result in rejection of your manuscript.

In accordance with BioMed Central editorial policies and formatting guidelines, all manuscript submissions to BMC Medicine must contain a Declarations section which includes the mandatory sub-sections listed below.

Please refer to the journal's Submission Guidelines web page for information regarding the criteria for each sub-section (https://bmcmedicine.biomedcentral.com/).

Where a mandatory Declarations section is not relevant to your study design or article type, please write 'Not applicable' in these sections.

For the 'Availability of data and materials' section, please provide information about

where the data supporting your findings can be found.

We encourage authors to deposit their datasets in publicly available repositories (where available and appropriate), or to be presented within the manuscript and/or additional supporting files.

Please note that identifying/confidential patient data should not be shared.

Authors who do not wish to share their data must confirm this under this sub-heading and also provide their reasons.

For further guidance on how to format this section, please refer to BioMed Central's editorial policies page (see links below).

Declarations Ethics approval and consent to participate Consent to publish Availability of data and materials -

Competing interests

-

Funding

_

Authors' Contributions

-

Acknowledgements

Further information about our editorial policies can be found at the following links: Ethical approval and consent:

http://www.biomedcentral.com/about/editorialpolicies#Ethics

Availability of data and materials section:

http://www.biomedcentral.com/submissions/editorial-

policies#availability+of+data+and+materials

If you need more time at any stage of the peer-review process, please do let us know. While our systems will continue to remind you of the original timelines, we aim to be as flexible as possible during the current pandemic.

This letter contains confidential information, is for your own use, and should not be

^{**}Our flexible approach during the COVID-19 pandemic**

forwarded to third parties.

Recipients of this email are registered users within the Editorial Manager database for this journal. We will keep your information on file to use in the process of submitting, evaluating and publishing a manuscript. For more information on how we use your personal details please see our privacy policy

at https://www.springernature.com/production-privacy-policy. If you no longer wish to receive messages from this journal or you have questions regarding database management, please contact the Publication Office at the link below.

Additional_file_1

Click here to access/download **Supplementary Material** Additional_file_1.docx

Additional_file_2

Click here to access/download **Supplementary Material** Additional_file_2.docx

Additional_file_3

Click here to access/download **Supplementary Material** Additional_file_3.docx